

NLRP3 Knockout 293T Cell Lysate, Homozygous

Catalog No.: RM02062

Basic Information

Catalog No.

RM02062

Category

Cell Lysate

Parental Cell line

293T

Genotype

Knockout

Gene Information

Gene Symbol

NLRP3

Species

Human

Gene ID

114548

Swiss Prot

Q96P20

SynonymsAGTAVPRL; All; AVP; C1orf7; CIAS1;
CLR1.1; FCAS; FCAS1; FCU; MWS; NALP3;
PYPAF1

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

This gene encodes a pyrin-like protein containing a pyrin domain, a nucleotide-binding site (NBS) domain, and a leucine-rich repeat (LRR) motif. This protein interacts with the apoptosis-associated speck-like protein PYCARD/ASC, which contains a caspase recruitment domain, and is a member of the NALP3 inflammasome complex. This complex functions as an upstream activator of NF-kappaB signaling, and it plays a role in the regulation of inflammation, the immune response, and apoptosis. Mutations in this gene are associated with familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome, and neonatal-onset multisystem inflammatory disease (NOMID). Multiple alternatively spliced transcript variants encoding distinct isoforms have been identified for this gene. Alternative 5' UTR structures are suggested by available data; however, insufficient evidence is available to determine if all of the represented 5' UTR splice patterns are biologically valid. [provided by RefSeq, Oct 2008]

Product Information

Description

NLRP3 Knockout 293T Cell Line is engineered from 293T cell line with Gene-Editing technology.

Allele-1:64bp deletion in exon1

Allele-2:85bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50µL, 2µg/µL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT GATCTAGCCACGCT*****AACAGGAGAGACCT
Mut GATCTAGCCACGCT***Deletion***AACAGGAGAGACCT
Allele-1: 64bp deletion in exon1
WT TGGATCTAGCCACG*****AGAAAGCAAAAAGA
Mut TGGATCTAGCCACG***Deletion***AGAAAGCAAAAAGA
Allele-2: 85bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and NLRP3 knockout (KO) 293T cells, using sanger sequencing.