

# NFKBIA Knockout HeLa Cell Lysate, Homozygous

Catalog No.: RM02080

### **Basic Information**

### Catalog No.

RM02080

### Category

Cell Lysate

### **Parental Cell line**

HeLa

### Genotype

Knockout

### **Background**

This gene encodes a member of the NF-kappa-B inhibitor family, which contain multiple ankrin repeat domains. The encoded protein interacts with REL dimers to inhibit NF-kappa-B/REL complexes which are involved in inflammatory responses. The encoded protein moves between the cytoplasm and the nucleus via a nuclear localization signal and CRM1-mediated nuclear export. Mutations in this gene have been found in ectodermal dysplasia anhidrotic with T-cell immunodeficiency autosomal dominant disease. [provided by RefSeq, Aug 2011]

### **Gene Information**

### **Gene Symbol**

**NFKBIA** 

### **Species**

Human

### Gene ID

4792

### **Swiss Prot**

P25963

### **Synonyms**

IKBA; MAD-3; NFKBI

### **Contact**

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	Τ	www.abclonal.com.cn

### **Product Information**

#### Description

NFKBIA Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.

Allele-1:77bp deletion in exon1

Allele-2:77bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

### **Packaging**

 ${\bf 1}$  vial parental cell Lysate and  ${\bf 1}$  vial knockout cell Lysate

Shipping Conditions
4°C
Amount
50μL, 2μg/μL.

### Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

#### Protocol

To be used as WB control. Lysate is supplied in  $1\times$  SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3-5 minutes before loading onto gel.

## Sequencing data

WT GACGACCGCCACGA\*\*\*\*\*\*\*\*AGGAGGTGCCGCGC
Mut GACGACCGCCACGA\*\*\*Deletion\*\*\*AGGAGGTGCCGCGC
Allele-1: 77bp deletion in exon1

WT GACGACCGCCACGA\*\*\*\*\*\*\*\*\*AGGAGGTGCCGCGC
Mut GACGACCGCCACGA\*\*\*Deletion\*\*\*AGGAGGTGCCGCGC
Allele-2: 77bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and NFKBIA knockout (KO) HeLa cells, using sanger sequencing.