

PER2 Knockout 293T Cell Line, Homozygous

Catalog No.: RM02103

Basic Information

Catalog No.

RM02103

Category

Cell Line

Parental Cell line

293T

Genotype

Knockout

Gene Information

Gene Symbol

PER2

Species

Human

Gene ID

8864

Swiss Prot

O15055

Synonyms

FASPS; FASPS1

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

This gene is a member of the Period family of genes and is expressed in a circadian pattern in the suprachiasmatic nucleus, the primary circadian pacemaker in the mammalian brain. Genes in this family encode components of the circadian rhythms of locomotor activity, metabolism, and behavior. This gene is upregulated by CLOCK/ARNTL heterodimers but then represses this upregulation in a feedback loop using PER/CRY heterodimers to interact with CLOCK/ARNTL. Polymorphisms in this gene may increase the risk of getting certain cancers and have been linked to sleep disorders. [provided by RefSeq, Jan 2014]

Product Information

Description

PER2 Knockout 293T Cell Line is engineered from 293T cell line with Gene-Editing Technology.

Allele-1:104bp deletion in exon1

Allele-2:104bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions

Dry ice

Amount

1~5x10⁶ cells/vial

Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

Protocol

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO₂ condition.

1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO₂.
7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

WT CCCCACCAAGGAGC*****TCGCAGGGCAGTGA
Mut CCCCACCAAGGAGC***Deletion***TCGCAGGGCAGTGA
Allele-1: 104bp deletion in exon1
WT CCCCACCAAGGAGC*****TCGCAGGGCAGTGA
Mut CCCCACCAAGGAGC***Deletion***TCGCAGGGCAGTGA
Allele-2: 104bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and PER2 knockout (KO) 293T cells, using sanger sequencing.