

Ndrg2 Knockout MA1800 Cell Line, Homozygous

Catalog No.: RM02110

Basic Information

Catalog No.

RM02110

Category

Cell Line

Parental Cell line

MA1800

Genotype

Knockout

Gene Information

Gene Symbol

Ndrg2

Species

Mouse

Gene ID

29811

Synonyms

AI182517; AU040374; Ndr2; SYLD

Contact

<u>a</u>	400-	999-6126
\bowtie	cn.market@abclona	al.com.cn
•	www.abclona	al.com.cn

Background

Product Information

Description

Ndrg2 Knockout MA1800 Cell Line is engineered from MA1800 cell line with Gene-Editing Technology.

Allele-1:1bp deletion in exon2

Allele-2:1bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

 ${\bf 1}$ vial parental cell line and ${\bf 1}$ vial knockout cell line

Shipping Conditions

Amount

Dry ice

1~5x10⁶ cells/vial

Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

Protoco

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37° C with 5% CO₂ condition.

- 1. Thaw the vial in 37°C water bath ,and shake it to melt as soon as possible.
- 2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
- 3. Remove and discard the supernatant.
- 4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
- 5. Add 8-10mL of complete medium.
- 6. Incubate the culture at 37°C incubator with 5% CO₂.
- 7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

WT TCTGTGGAGACACCTTATGGCTCGGTCACTTTTACTGTG
Mut TCTGTGGAGACACCTTATG -CTCGGTCACTTTTACTGTG

Allele-1: 1bp deletion in exon2

WT CCAAACCCAAACGCCCGGCGATATTCACCTACCATGATG
Mut CCAAACCCAAACGCCCGGC -ATATTCACCTACCATGATG

Allele-2: 1bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and Ndrg2 knockout (KO) MA1800 cells, using sanger sequencing.