

Src Knockout HeLa Cell Line, Homozygous

Catalog No.: RM02114

Basic Information

Catalog No.

RM02114

Category

Cell Line

Parental Cell line

HeLa

Genotype

Knockout

Gene Information

Gene Symbol

Src

Species

Human

Gene ID

6714

Swiss Prot

P12931

Synonyms

ASV; SRC1; THC6; c-SRC; p60-Src

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

This gene is highly similar to the v-src gene of Rous sarcoma virus. This proto-oncogene may play a role in the regulation of embryonic development and cell growth. The protein encoded by this gene is a tyrosine-protein kinase whose activity can be inhibited by phosphorylation by c-SRC kinase. Mutations in this gene could be involved in the malignant progression of colon cancer. Two transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul 2008]

Product Information

Description

Src Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing Technology.
Allele-1:46bp deletion in exon2
Allele-2:46bp deletion in exon2
Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions

Dry ice

Amount

1~5x10⁶ cells/vial

Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

Protocol

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO₂ condition.

1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO₂.
7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

WT TGACTATGAGTCTA*****TGCAACAACACGT
Mut TGACTATGAGTCTA***Deletion***TGCAACAACACGT
Allele-1: 46bp deletion in exon2
WT TGACTATGAGTCTA*****TGCAACAACACGT
Mut TGACTATGAGTCTA***Deletion***TGCAACAACACGT
Allele-2: 46bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and Src knockout (KO) HeLa cells, using sanger sequencing.