

CD63 Knockout 293T Cell Lysate, Homozygous

Catalog No.: RM02188

Basic Information

Catalog No.

RM02188

Category

Cell Lysate

Parental Cell line

293T

Genotype

Knockout

Gene Information

Gene Symbol

CD63

Species

Human

Gene ID

967

Swiss Prot

P08962

SynonymsLAMP-3; ME491; MLA1; OMA81H;
TSPAN30

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. The encoded protein is a cell surface glycoprotein that is known to complex with integrins. It may function as a blood platelet activation marker. Deficiency of this protein is associated with Hermansky-Pudlak syndrome. Also this gene has been associated with tumor progression. Alternative splicing results in multiple transcript variants encoding different protein isoforms. [provided by RefSeq, Apr 2012]

Product Information

Description

CD63 Knockout 293T Cell Line is engineered from 293T cell line with Gene-Editing technology.

Allele-1:77bp deletion in exon3

Allele-2:77bp deletion in exon3

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50µL, 2µg/µL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol).

Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT ATCCCTCCATGTC*****ATAAGGAGGTAGGG
Mut ATCCCTCCATGTC***Deletion***ATAAGGAGGTAGGG
Allele-1: 77bp deletion in exon3
WT ATCCCTCCATGTC*****ATAAGGAGGTAGGG
Mut ATCCCTCCATGTC***Deletion***ATAAGGAGGTAGGG
Allele-2: 77bp deletion in exon3

Genome sequence analysis of PCR products from parental (WT) and CD63 knockout (KO) 293T cells, using sanger sequencing.