

MYD88 Knockout HeLa Cell Lysate, Homozygous

Catalog No.: RM02192

Basic Information

Catalog No.

RM02192

Category

Cell Lysate

Parental Cell line

HeLa

Genotype

Knockout

Gene Information

Gene Symbol

MYD88

Species

Human

Gene ID

4615

Swiss Prot

Q99836

Synonyms

MYD88D

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

This gene encodes a cytosolic adapter protein that plays a central role in the innate and adaptive immune response. This protein functions as an essential signal transducer in the interleukin-1 and Toll-like receptor signaling pathways. These pathways regulate that activation of numerous proinflammatory genes. The encoded protein consists of an N-terminal death domain and a C-terminal Toll-interleukin1 receptor domain. Patients with defects in this gene have an increased susceptibility to pyogenic bacterial infections. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Feb 2010]

Product Information

Description

MYD88 Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.

Allele-1:65bp deletion in exon1

Allele-2:85bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50μL, 2μg/μL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT CTTGAACGTGCGGA*****CAACTGGAGACACA
Mut CTTGAACGTGCGGA***Deletion***AACTGGAGACACA
Allele-1: 65bp deletion in exon1

WT CTTGAACGTGCGGA*****CCCCACTGGCAGGC
Mut CTTGAACGTGCGGA***Deletion***CCCCACTGGCAGGC
Allele-2: 85bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and MYD88 Knockout (KO) HeLa cells, using sanger sequencing.