

# BRD4 Knockout HeLa Cell Line, Homozygous

Catalog No.: RM02204

## Basic Information

**Catalog No.**

RM02204

**Category**

Cell Line

**Parental Cell line**

HeLa

**Genotype**

Knockout

## Gene Information

**Gene Symbol**

BRD4

**Species**

Human

**Gene ID**

23476

**Swiss Prot**

O60885

**Synonyms**

CAP; HUNK1; HUNKI; MCAP

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

The protein encoded by this gene is homologous to the murine protein MCAP, which associates with chromosomes during mitosis, and to the human RING3 protein, a serine/threonine kinase. Each of these proteins contains two bromodomains, a conserved sequence motif which may be involved in chromatin targeting. This gene has been implicated as the chromosome 19 target of translocation t(15;19)(q13;p13.1), which defines an upper respiratory tract carcinoma in young people. Two alternatively spliced transcript variants have been described. [provided by RefSeq, Jul 2008]

## Product Information

**Description**

BRD4 Knockout cell line is engineered from HeLa cell line with Gene-Editing Technology.

Allele-1:73bp deletion in exon1

Allele-2:73bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

**Packaging**

1 vial parental cell line and 1 vial knockout cell line

**Shipping Conditions**

Dry ice

**Amount**

1~5x10<sup>6</sup> cells/vial

**Storage**

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

**Protocol**

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO<sub>2</sub> condition.

1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO<sub>2</sub>.
7. A subcultivation ratio of 1:2-1:4 is recommended.

## Sequencing data

---

WT AGACCAACCAACTG\*\*\*\*\*ATGCCGTCAAGCTG  
Mut AGACCAACCAACTG\*\*\*Deletion\*\*\*ATGCCGTCAAGCTG  
Allele-1: 73bp deletion in exon1  
WT AGACCAACCAACTG\*\*\*\*\*ATGCCGTCAAGCTG  
Mut AGACCAACCAACTG\*\*\*Deletion\*\*\*ATGCCGTCAAGCTG  
Allele-2: 73bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and BRD4 knockout (KO) HeLa cells, using sanger sequencing.