Leader in Biomolecular Solutions for Life Science

www.abclonal.com

HK2 Knockout HeLa Cell Line, Homozygous

Catalog No.: RM02205

Basic Information

Catalog No.
RM02205

Category
Cell Line

Parental Cell line
HeLa

Genotype
Knockout

Background

Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most glucose metabolism pathways. This gene encodes hexokinase 2, the predominant form found in skeletal muscle. It localizes to the outer membrane of mitochondria. Expression of this gene is insulin-responsive, and studies in rat suggest that it is involved in the increased rate of glycolysis seen in rapidly growing cancer cells. [provided by RefSeq, Apr 2009]

Gene Information

Gene Symbol
HK2

Species
Human

Gene ID
3099

Swiss Prot
P52789

Synonyms
HKII; HXK2

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Product Information

Description
HK2 Knockout cell line is engineered from HeLa cell line with Gene-Editing Technology.
Allele-1:exon2 was deleted
Allele-2:exon2 was deleted
Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging
1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions
Dry ice

Amount
1~5x10^6 cells/vial

Storage
Store at -20℃

Protocol
Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37℃ with 5% CO_2 condition.

1. Thaw the vial in 37℃ water bath ,and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37℃ incubator with 5% CO_2.
7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

WT GTGGTGTTCCATGA**********CTCAGGGGTGATTT
Mut GTGGTGTTCCATGA***Deletion***CTCAGGGGTGATTT
Allele-1: exon2 was deleted

WT TGAAGAGCTGAGTG**********CTCAGGGGTGATTT
Mut TGAAGAGCTGAGTC***Deletion***CTCAGGGGTGATTT
Allele-2: exon2 was deleted

Genome sequence analysis of PCR products from parental (WT) and HK2 knockout (KO) HeLa cells, using sanger sequencing.

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.