

ANXA1 Knockdown HeLa Cell Line, Heterozygous

Catalog No.: RM02225 **1 Publications**

Basic Information

Catalog No.

RM02225

Category

Cell Line

Parental Cell line

NIH/3T3

Genotype

Knockdown

Gene Information

Gene Symbol

ANXA1

Species

Human

Gene ID

301

Swiss Prot

P04083

Synonyms

ANX1; LPC1

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

This gene encodes a membrane-localized protein that binds phospholipids. This protein inhibits phospholipase A2 and has anti-inflammatory activity. Loss of function or expression of this gene has been detected in multiple tumors. [provided by RefSeq, Dec 2014]

Product Information

Description

ANXA1 Knockdown HeLa Cell Line is engineered from NIH/3T3 cell line with Gene-Editing Technology.

Allele-1:53bp deletion in exon2

Allele-2:54bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions

Dry ice

Amount

1~5x10⁶ cells/vial

Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

Protocol

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO₂ condition.

1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO₂.
7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

WT AGGTGGTCCCGGAT*****CATAAGGCCATAAT
Mut AGGTGGTCCCGGAT***Deletion***CATAAGGCCATAAT
Allele-1: 53bp deletion in exon2
WT AGGTGGTCCCGGAT*****ATAAGGCCATAATG
Mut AGGTGGTCCCGGAT***Deletion***ATAAGGCCATAATG
Allele-2: 54bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and ANXA1 Knockdown (KD) HeLa cells, using sanger sequencing.