# ABclonal www.abclonal.com

# **CETN2 Knockout HeLa Cell Lysate, Homozygous**

Catalog No.: RM02297

#### **Basic Information**

#### Catalog No.

RM02297

#### Category

Cell Lysate

# **Parental Cell line**

HeLa

#### Genotype

Knockout

# **Background**

Caltractin belongs to a family of calcium-binding proteins and is a structural component of the centrosome. The high level of conservation from algae to humans and its association with the centrosome suggested that caltractin plays a fundamental role in the structure and function of the microtubule-organizing center, possibly required for the proper duplication and segregation of the centrosome. [provided by RefSeq, Jul 2008]

#### **Gene Information**

#### **Gene Symbol**

CETN2

## **Species**

Human

#### **Gene ID**

1069

#### **Swiss Prot**

P41208

# Synonyms

CALT; CEN2

#### **Contact**

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	Τ	www.abclonal.com.cn

#### **Product Information**

#### **Description**

CETN2 Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.

Allele-1:80bp deletion in exon1

Allele-2:80bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

## **Packaging**

1 vial parental cell Lysate and 1 vial knockout cell Lysate

#### **Shipping Conditions**

Amount

4°C

50μL, 2μg/μL.

#### Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

#### **Protocol**

To be used as WB control. Lysate is supplied in  $1\times$  SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

# Sequencing data

WT AGAATGAGCCCTAA\*\*\*\*\*\*\*\*\*\*\*ATGTTAAAGAACTG
Mut AGAATGAGCCCTAA\*\*\*Deletion\*\*\*ATGTTAAAGAACTG
Allele-1: 80bp deletion in exon1

WT AGAATGAGCCCTAA\*\*\*\*\*\*\*ATGTTAAAGAACTG
Mut AGAATGAGCCCTAA\*\*\*Deletion\*\*\*ATGTTAAAGAACTG

Allele-2: 80bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and CETN2 Knockout (KO) HeLa cells, using sanger sequencing.