

TDO2 Knockdown 293T Cell Lysate, Heterozygous

Catalog No.: RM02300

Basic Information

Catalog No.

RM02300

Category

Cell Lysate

Parental Cell line

293T

Genotype

Knockdown

Gene Information

Gene Symbol

TDO2

Species

Human

Gene ID

6999

Swiss Prot

P48775

Synonyms

TDO; TO; TPH2; TRPO

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

This gene encodes a heme enzyme that plays a critical role in tryptophan metabolism by catalyzing the first and rate-limiting step of the kynurenine pathway. Increased activity of the encoded protein and subsequent kynurenine production may also play a role in cancer through the suppression of antitumor immune responses, and single nucleotide polymorphisms in this gene may be associated with autism. [provided by RefSeq, Feb 2012]

Product Information

Description

TDO2 Knockdown 293T Cell Line is engineered from 293T cell line with Gene-Editing technology.

Allele-1:62bp deletion in exon2

Allele-2:63bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50µL, 2µg/µL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT AAAAACTCCCGTA*****CTATGGGA ACTACC
Mut AAAAACTCCCGTA***Deletion***TATGGGA ACTACC
Allele-1: 62bp deletion in exon2
WT AAAAACTCCCGTA*****TATGGGA ACTACCT
Mut AAAAACTCCCGTA***Deletion***TATGGGA ACTACCT
Allele-2: 63bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and TDO2 Knockdown (KD) 293T cells, using sanger sequencing.