

TGFB2 Knockdown 293T Cell Lysate, Heterozygous

Catalog No.: RM02307

Basic Information

Catalog No.

RM02307

Category

Cell Lysate

Parental Cell line

293T

Genotype

Knockdown

Gene Information

Gene Symbol

TGFB2

Species

Human

Gene ID

7048

Swiss Prot

P37173

SynonymsAAT3; FAA3; LDS1B; LDS2; LDS2B; MFS2;
RIIC; TAAD2; TGFR-2; TGFbeta-RII

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

This gene encodes a member of the Ser/Thr protein kinase family and the TGFB receptor subfamily. The encoded protein is a transmembrane protein that has a protein kinase domain, forms a heterodimeric complex with another receptor protein, and binds TGF-beta. This receptor/ligand complex phosphorylates proteins, which then enter the nucleus and regulate the transcription of a subset of genes related to cell proliferation. Mutations in this gene have been associated with Marfan Syndrome, Loeys-Deitz Aortic Aneurysm Syndrome, and the development of various types of tumors. Alternatively spliced transcript variants encoding different isoforms have been characterized. [provided by RefSeq, Jul 2008]

Product Information

Description

TGFB2 Knockdown 293T Cell Line is engineered from 293T cell line with Gene-Editing technology.

Allele-1:21bp deletion and 8bp deletion in exon3

Allele-2:90bp deletion in exon3

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50μL, 2μg/μL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT CAGTTTGCCATGAC*****TTTATTCTGGGAAGA***CTTCATGTGTCTCTGAGCTCTGATGAGTGCAATGA
Mut CAGTTTGCCATGAC***Deletion***TTTATTCTGGGAAGA***CTTCATGTGTCTCT - - - - - GATGAGTGCAATGA
Allele-1: 21bp deletion and 8bp deletion in exon3
WT CAAGCTCCCCCTACC*****GCTCTGATGAGTGC
Mut CAAGCTCCCCCTACC***Deletion***GCTCTGATGAGTGC
Allele-2: 90bp deletion in exon3

Genome sequence analysis of PCR products from parental (WT) and TGFBR2 Knockdown (KD) 293T cells, using sanger sequencing.