

## CDKL4 Knockout NIH/3T3 Cell Lysate, Homozygous

Catalog No.: RM02317

## **Basic Information**

## Catalog No.

RM02317

## Category

Cell Lysate

## **Parental Cell line**

NIH/3T3

#### Genotype

Knockout

# Gene Information

## **Gene Symbol**

CDKL4

## **Species**

Mouse

#### **Gene ID**

381113

#### **Synonyms**

AU067824; Gm942

## **Contact**

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	Τ	www.abclonal.com.cn

## **Background**

## **Product Information**

#### **Description**

CDKL4 Knockout NIH/3T3 Cell Line is engineered from NIH/3T3 cell line with Gene-Editing technology.

Allele-1:61bp deletion in exon3

Allele-2:2bp deletion in exon3

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

#### **Packaging**

1 vial parental cell Lysate and 1 vial knockout cell Lysate

**Shipping Conditions** 4°C

**Amount** 50μL, 2μg/μL.

#### Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

#### **Protocol**

To be used as WB control. Lysate is supplied in  $1\times$  SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

## Sequencing data

WT AGTTGAAACACCCA\*\*\*\*\*\*\*\*\*\*\*\*CCCAAACGGGTAAG
Mut AGTTGAAACACCCA\*\*\*Deletion\*\*\*CCCAAACGGGTAAG
Allele-1: 61bp deletion in exon3

WT CCCAAACCTCGTGA\*\*\*\*\*\*\*\*\*\*GTTAAACGAGCTGG
Mut CCCAAAC\_TCGTGA\*\*\*\*\*\*\*\*GTTAAAC\_AGCTGG
Allele-2: 2bp deletion in exon3

Genome sequence analysis of PCR products from parental (WT) and CDKL4 knockout (KO) NIH3T3 cells, using sanger sequencing.