

# TLK1 Knockdown NIH/3T3 Cell Lysate, Heterozygous

Catalog No.: RM02333

# **Basic Information**

# Catalog No.

RM02333

## Category

Cell Lysate

### **Parental Cell line**

NIH/3T3

#### Genotype

Knockdown

# Gene Information

#### **Gene Symbol**

TLK1

# **Species**

Mouse

#### **Gene ID**

228012

#### **Synonyms**

4930545J15Rik

# **Contact**

<u>a</u>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

# **Background**

## **Product Information**

#### **Description**

TLK1 Knockdown NIH/3T3 Cell Line is engineered from NIH/3T3 cell line with Gene-Editing technology.

Allele-1:43bp deletion in exon1

Allele-2:43bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

#### **Packaging**

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping ConditionsAmount $4^{\circ}C$  $50\mu L$ ,  $2\mu g/\mu L$ .

#### Storage

Lysate is stable for 12 months when stored at -20  $^{\circ}\text{C}.$  Minimizing freeze-thaw cycles.

#### **Protocol**

To be used as WB control. Lysate is supplied in  $1\times$  SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

# Sequencing data

WT CCAACCCCGGGCTC\*GGAGGCCCAGGGA
Mut CCAACCCCGGGCTC\*\*\*Deletion\*\*\*\*GGGAGGCCCAGGGA

Allele-1: 43bp deletion in exon1

WT CTCCAACCCCGGGC\*\*\*\*\*\*\*GGGAGGCCCAGGGA
Mut CTCCAACCCCGGGC\*\*\*Deletion\*\*\*GGGAGGCCCAGGGA

Allele-2: 43bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and TLK1 Knockdown (KD) NIH3T3 cells, using sanger sequencing.