

E2F1 Knockout HeLa Cell Lysate, Homozygous

Catalog No.: RM02341

Basic Information

Catalog No.

RM02341

Category

Cell Lysate

Parental Cell line

HeLa

Genotype

Knockout

Gene Information

Gene Symbol

E2F1

Species

Human

Gene ID

1869

Swiss Prot

Q01094

Synonyms

E2F-1; RBAP1; RBBP3; RBP3

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

The protein encoded by this gene is a member of the E2F family of transcription factors. The E2F family plays a crucial role in the control of cell cycle and action of tumor suppressor proteins and is also a target of the transforming proteins of small DNA tumor viruses. The E2F proteins contain several evolutionally conserved domains found in most members of the family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor proteins (DP), a transactivation domain enriched in acidic amino acids, and a tumor suppressor protein association domain which is embedded within the transactivation domain. This protein and another 2 members, E2F2 and E2F3, have an additional cyclin binding domain. This protein binds preferentially to retinoblastoma protein pRB in a cell-cycle dependent manner. It can mediate both cell proliferation and p53-dependent/independent apoptosis. [provided by RefSeq, Jul 2008]

Product Information

Description

E2F1 Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.

Allele-1:79bp deletion in exon3

Allele-2:79bp deletion in exon3

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50μL, 2μg/μL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT TGCTGAGCCACTCG*****TCCTTGAGGGCATC
Mut TGCTGAGCCACTCG***Deletion***TCCTTGAGGGCATC
Allele-1: 79bp deletion in exon3
WT TGCTGAGCCACTCG*****TCCTTGAGGGCATC
Mut TGCTGAGCCACTCG***Deletion***TCCTTGAGGGCATC
Allele-2: 79bp deletion in exon3

Genome sequence analysis of PCR products from parental (WT) and E2F1 Knockout (KO) HeLa cells, using sanger sequencing.