

COX4I1 Knockdown 293T Cell Lysate, Heterozygous

Catalog No.: RM02354

Basic Information

Catalog No.

RM02354

Category

Cell Lysate

Parental Cell line

293T

Genotype

Knockdown

Gene Information

Gene Symbol

COX4I1

Species

Human

Gene ID

1327

Swiss Prot

P13073

SynonymsCOX IV-1; COX4; COX4-1; COXIV;
COXIV-1

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

Cytochrome c oxidase (COX) is the terminal enzyme of the mitochondrial respiratory chain. It is a multi-subunit enzyme complex that couples the transfer of electrons from cytochrome c to molecular oxygen and contributes to a proton electrochemical gradient across the inner mitochondrial membrane. The complex consists of 13 mitochondrial- and nuclear-encoded subunits. The mitochondrially-encoded subunits perform the electron transfer and proton pumping activities. The functions of the nuclear-encoded subunits are unknown but they may play a role in the regulation and assembly of the complex. This gene encodes the nuclear-encoded subunit IV isoform 1 of the human mitochondrial respiratory chain enzyme. It is located at the 3' of the NOC4 (neighbor of COX4) gene in a head-to-head orientation, and shares a promoter with it. Pseudogenes related to this gene are located on chromosomes 13 and 14. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jan 2016]

Product Information

Description

COX4I1 Knockdown 293T Cell Line is engineered from 293T cell line with Gene-Editing technology.

Allele-1:29bp deletion and 1bp deletion in exon2

Allele-2:65bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50µL, 2µg/µL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT TTTTCGCTCCCAGC*****CGGAGGTGGCCCAT***CACCTGTCTGCCAGCCAGAAGGCACTGAAG
Mut TTTTCGCTCCCAGC***Deletion***CGGAGGTGGCCCAT***CACCTGTCTGCCAGC -AGAAGGCACTGAAG
Allele-1: 28bp deletion and 1bp deletion in exon2
WT TTTTCGCTCCCAGC*****AGAAGGCACTGAAG
Mut TTTTCGCTCCCAGC***Deletion***AGAAGGCACTGAAG
Allele-2: 65bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and COX4I1 Knockdown (KD) 293T cells, using sanger sequencing.