

# CAST Knockout HeLa Cell Lysate, Homozygous

Catalog No.: RM02374

# **Basic Information**

#### Catalog No.

RM02374

# Category

Cell Lysate

#### **Parental Cell line**

HeLa

#### Genotype

Knockout

# **Background**

The protein encoded by this gene is an endogenous calpain (calcium-dependent cysteine protease) inhibitor. It consists of an N-terminal domain L and four repetitive calpain-inhibition domains (domains 1-4), and it is involved in the proteolysis of amyloid precursor protein. The calpain/calpastatin system is involved in numerous membrane fusion events, such as neural vesicle exocytosis and platelet and red-cell aggregation. The encoded protein is also thought to affect the expression levels of genes encoding structural or regulatory proteins. Alternatively spliced transcript variants encoding different isoforms have been described. [provided by RefSeq, Jun 2010]

#### **Gene Information**

## **Gene Symbol**

CAST

## **Species**

Human

# **Gene ID**

831

#### **Swiss Prot**

P20810

# Synonyms

BS-17; PLACK

## **Contact**

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

# **Product Information**

#### **Description**

CAST Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.

Allele-1:1bp deletion in exon4

Allele-2:2bp deletion in exon4

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

# **Packaging**

1 vial parental cell Lysate and 1 vial knockout cell Lysate

## **Shipping Conditions**

Amount

4°C

50μL, 2μg/μL.

#### Storage

Lysate is stable for 12 months when stored at -20  $^{\circ}$ C. Minimizing freeze-thaw cycles.

#### **Protocol**

To be used as WB control. Lysate is supplied in  $1\times$  SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

# Sequencing data

WT CAAGCAGGCATCAGATACAGGAAGTAACG
Mut CAAGCAGGCATCAG-TACAGGAAGTAACG

Allele-1: 1bp deletion in exon4

WT CCAAGCAGGCATCAGATACAGGAAGTAACG
Mut CCAAGCAGGCATCA - -TACAGGAAGTAACG

Allele-2: 2bp deletion in exon4

Genome sequence analysis of PCR products from parental (WT) and CAST knockout (KO) HeLa cells, using sanger sequencing.