Leader in Biomolecular Solutions for Life Science

www.abclonal.com

FGFR2 Knockout 293T Cell Lysate, Homozygous

Catalog No.: RM02400

Basic Information

Catalog No.
RM02400

Category
Cell Lysate

Parental Cell line
293T

Genotype
Knockout

Gene Information

Gene Symbol
FGFR2

Species
Human

Gene ID
2263

Swiss Prot
P21802

Synonyms
BBDS; BEK; BFR-1; CD332; CEK3; CFD1; ECT1; JWS; K-SAM; KGFR; TK14; TK25

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Background

The protein encoded by this gene is a member of the fibroblast growth factor receptor family, where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein consists of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. This particular family member is a high-affinity receptor for acidic, basic and/or keratinocyte growth factor, depending on the isoform. Mutations in this gene are associated with Crouzon syndrome, Pfeiffer syndrome, Craniosynostosis, Apert syndrome, Jackson-Weiss syndrome, Beare-Stevenson cutis gyrata syndrome, Saethre-Chotzen syndrome, and syndromic craniosynostosis. Multiple alternatively spliced transcript variants encoding different isoforms have been noted for this gene. [provided by RefSeq, Jan 2009]

Product Information

Description
FGFR2 Knockout 293T Cell Line is engineered from 293T cell line with Gene-Editing technology.
Allele-1:exon8 was deleted
Allele-2:exon8 was deleted
Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging
1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions
4℃

Amount
50μL, 2μg/μL.

Storage
Store at -20℃

Protocol
To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

```
WT    CCCCATCACCAGAT**************AGCGGGGAGGCGGG
Mut   CCCCATCACCAGAT***Deletion***AGCGGGGAGGCGGG
Allele-1: exon8 was deleted

WT    CCCCATCACCAGAT**************AGCGGGGAGGCGGG
Mut   CCCCATCACCAGAT***Deletion***AGCGGGGAGGCGGG
Allele-2: exon8 was deleted
```

Genome sequence analysis of PCR products from parental (WT) and FGFR2 knockout (KO) 293T cells, using sanger sequencing.