

CTNND1 Knockout HeLa Cell Line, Homozygous

Catalog No.: RM02414

Basic Information

Catalog No.

RM02414

Category

Cell Line

Parental Cell line

HeLa

Genotype

Knockout

Gene Information

Gene Symbol

CTNND1

Species

Human

Gene ID

1500

Swiss Prot

O60716

Synonyms

CAS; CTNND; P120CAS; P120CTN; p120; p120(CAS); p120(CTN)

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

This gene encodes a member of the Armadillo protein family, which function in adhesion between cells and signal transduction. Multiple translation initiation codons and alternative splicing result in many different isoforms being translated. Not all of the full-length natures of the described transcript variants have been determined. Read-through transcription also exists between this gene and the neighboring upstream thioredoxin-related transmembrane protein 2 (TMX2) gene. [provided by RefSeq, Dec 2010]

Product Information

Description

CTNND1 Knockout HeLa Cell Line knockout is engineered from HeLa cell line with Gene-Editing Technology.

Allele-1:106bp deletion in exon3

Allele-2:106bp deletion in exon3

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions

Dry ice

Amount

1~5x10⁶ cells/vial

Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

Protocol

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO₂ condition.

1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO₂.
7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

WT TATATAGCACCATC*****GGCGCACAGAGACC
Mut TATATAGCACCATC***Deletion***GGCGCACAGAGACC
Allele-1: 106bp deletion in exon3
WT TATATAGCACCATC*****GGCGCACAGAGACC
Mut TATATAGCACCATC***Deletion***GGCGCACAGAGACC
Allele-2: 106bp deletion in exon3

Genome sequence analysis of PCR products from parental (WT) and CTNND1 knockout (KO) HeLa cells, using sanger sequencing.