

TOP2A Knockout 293T Cell Lysate, Homozygous

Catalog No.: RM02473

Basic Information

Catalog No.

RM02473

Category

Cell Lysate

Parental Cell line

293T

Genotype

Knockout

Gene Information

Gene Symbol

TOP2A

Species

Human

Gene ID

7153


Swiss Prot

P11388

Synonyms

TOP2; TP2A

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

This gene encodes a DNA topoisomerase, an enzyme that controls and alters the topologic states of DNA during transcription. This nuclear enzyme is involved in processes such as chromosome condensation, chromatid separation, and the relief of torsional stress that occurs during DNA transcription and replication. It catalyzes the transient breaking and rejoining of two strands of duplex DNA which allows the strands to pass through one another, thus altering the topology of DNA. Two forms of this enzyme exist as likely products of a gene duplication event. The gene encoding this form, alpha, is localized to chromosome 17 and the beta gene is localized to chromosome 3. The gene encoding this enzyme functions as the target for several anticancer agents and a variety of mutations in this gene have been associated with the development of drug resistance. Reduced activity of this enzyme may also play a role in ataxia-telangiectasia. [provided by RefSeq, Jul 2010]

Product Information

Description

TOP2A Knockout 293T Cell Line is engineered from 293T cell line with Gene-Editing technology.

Allele-1:97bp deletion in exon1

Allele-2:97bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50µL, 2µg/µL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT CTTGAGCCCCTTCA*****GTGACGGGTGAAGC
Mut CTTGAGCCCCTTCA***Deletion***GTGACGGGTGAAGC
Allele-1: 97bp deletion in exon1
WT CTTGAGCCCCTTCA*****GTGACGGGTGAAGC
Mut CTTGAGCCCCTTCA***Deletion***GTGACGGGTGAAGC
Allele-2: 97bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and TOP2A knockout (KO) 293T cells, using sanger sequencing.