

# IGF1 Knockdown HeLa Cell Lysate, Heterozygous

Catalog No.: RM02550

### **Basic Information**

#### Catalog No.

RM02550

### Category

Cell Lysate

### **Parental Cell line**

HeLa

### Genotype

Knockdown

### **Background**

The protein encoded by this gene is similar to insulin in function and structure and is a member of a family of proteins involved in mediating growth and development. The encoded protein is processed from a precursor, bound by a specific receptor, and secreted. Defects in this gene are a cause of insulin-like growth factor I deficiency. Alternative splicing results in multiple transcript variants encoding different isoforms that may undergo similar processing to generate mature protein. [provided by RefSeq, Sep 2015]

### **Gene Information**

### **Gene Symbol**

IGF1

### **Species**

Human

### Gene ID

3479

#### **Swiss Prot**

P05019

### **Synonyms**

IGF-I; IGFI; MGF

### **Contact**

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	Τ	www.abclonal.com.cn

### **Product Information**

### **Description**

IGF1 Knockdown HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.

Allele-1:71bp deletion in exon2

Allele-2:WT

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

### **Packaging**

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping ConditionsAmount $4^{\circ}C$  $50\mu L$ ,  $2\mu g/\mu L$ .

#### Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

### **Protocol**

To be used as WB control. Lysate is supplied in  $1\times$  SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

## Sequencing data

WT CACCATGTCCTCCT\*\*\*\*\*\*\*\*\*\*TGCGGGGCTGAGCT
Mut CACCATGTCCTCCT\*\*\*Deletion\*\*\*TGCGGGGCTGAGCT
Allele-1: 71bp deletion in exon2

WT CTGGCGCTGTGCCTGCTCACCTTCACCAGCTCTGCCACGG Mut CTGGCGCTGTGCCTGCTCACCTTCACCAGCTCTGCCACGG Allele-2: WT Genome sequence analysis of PCR products from parental (WT) and IGF1 Knockdown (KD) HeLa cells, using sanger sequencing.