

IL18 Knockout HeLa Cell Lysate, Homozygous

Catalog No.: RM02560

Basic Information

Catalog No.

RM02560

Category

Cell Lysate

Parental Cell line

HeLa

Genotype

Knockout

Background

The protein encoded by this gene is a proinflammatory cytokine that augments natural killer cell activity in spleen cells, and stimulates interferon gamma production in T-helper type I cells. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Aug 2011]

Gene Information

Gene Symbol

IL18

Species

Human

Gene ID

3606

Swiss Prot

Q14116

Synonyms

IGIF; IL-18; IL-1g; IL1F4

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Product Information

Description

IL18 Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology. Allele-1:7bp deletion in exon2

Allele-2:52bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

Amount 50μL, 2μg/μL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in $1\times$ SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT CCAAGTTCTCTTCA*********AGGAAATCGGCCTC
Mut CCAAGTTCTCTTCA***Deletion***AGGAAATCGGCCTC
Allele-1: 7bp deletion in exon2

WT AAGTTCTCTTCATT**********TTTAATTCGCAAAC
Mut AAGTTCTCTTCATT***Deletion***TTTAATTCGCAAAC

Allele-2: 52bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and IL18 knockout (KO) HeLa cells, using sanger sequencing.