Leader in Biomolecular Solutions for Life Science

www.abclonal.com

# LRP6 Knockdown 293T Cell Lysate, Heterozygous

**Catalog No.: RM02564**

## Basic Information

**Catalog No.**
RM02564

**Category**
Cell Lysate

**Parental Cell line**
293T

**Genotype**
Knockdown

## Gene Information

**Gene Symbol**
LRP6

**Species**
Human

**Gene ID**
4040

**Swiss Prot**
O75581

**Synonyms**
ADCAD2; STHAG7

## Contact


400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn


## Background

This gene encodes a member of the low density lipoprotein (LDL) receptor gene family. LDL receptors are transmembrane cell surface proteins involved in receptor-mediated endocytosis of lipoprotein and protein ligands. The protein encoded by this gene functions as a receptor or, with Frizzled, a co-receptor for Wnt and thereby transmits the canonical Wnt/beta-catenin signaling cascade. Through its interaction with the Wnt/beta-catenin signaling cascade this gene plays a role in the regulation of cell differentiation, proliferation, and migration and the development of many cancer types. This protein undergoes gamma-secretase dependent RIP- (regulated intramembrane proteolysis) processing but the precise locations of the cleavage sites have not been determined.[provided by RefSeq, Dec 2009]

## Product Information

**Description**
LRP6 Knockdown 293T Cell Line is engineered from 293T cell line with Gene-Editing technology.
Allele-1:201bp deletion in exon2
Allele-2:199bp deletion in exon2
Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

**Packaging**
1 vial parental cell Lysate and 1 vial knockout cell Lysate

**Shipping Conditions**
4℃

**Amount**
50μL, 2μg/μL.

**Storage**
Store at -20℃

**Protocol**
To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

## Sequencing data

```
WT    AGACGGGACTTGCG**************GCTGGCATGTGATT
Mut   AGACGGGACTTGCG***Deletion***GCTGGCATGTGATT
Allele-1:    201bp deletion in exon2

WT    CAGACGGGACTTGC**************TGGGCTGGCATGTG
Mut   CAGACGGGACTTGC***Deletion***TGGGCTGGCATGTG
Allele-2:    199bp deletion in exon2
```

Genome sequence analysis of PCR products from parental (WT) and LRP6 Knockdown (KD) 293T cells, using sanger sequencing.