

CXCL8 Knockout HeLa Cell Line, Homozygous

Catalog No.: RM02622

Basic Information

Catalog No.

RM02622

Category

Cell Line

Parental Cell line

HeLa

Genotype

Knockout

Gene Information

Gene Symbol

CXCL8

Species

Human

Gene ID

3576

Swiss Prot

P10145

Synonyms

GCP-1; GCP1; IL8; LECT; LUCT; LYNAP;
MDNCF; MONAP; NAF; NAP-1; NAP1

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

The protein encoded by this gene is a member of the CXC chemokine family. This chemokine is one of the major mediators of the inflammatory response. This chemokine is secreted by several cell types. It functions as a chemoattractant, and is also a potent angiogenic factor. This gene is believed to play a role in the pathogenesis of bronchiolitis, a common respiratory tract disease caused by viral infection. This gene and other ten members of the CXC chemokine gene family form a chemokine gene cluster in a region mapped to chromosome 4q. [provided by RefSeq, Jul 2008]

Product Information

Description

CXCL8 Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.

Allele-1:53bp deletion in exon2

Allele-2:70bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions

Dry ice

Amount

1~5x10⁶ cells/vial

Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

Protocol

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO₂ condition.

1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO₂.
7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

WT GACATACTCAAAC*****TGCGCCAACACAGA
Mut GACATACTCAAAC***Deletion***TGCGCCAACACAGA
Allele-1: 53bp deletion in exon2

WT AGTGCATAAAGACA*****ACACAGAAATTATG
Mut AGTGCATAAAGACA***Deletion***ACACAGAAATTATG
Allele-2: 70bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and CXCL8 knockout (KO) HeLa cells, using sanger sequencing.