

SLC38A2 Knockout HeLa Cell Line, Homozygous

Catalog No.: RM02679

Basic Information

Catalog No.

RM02679

Category

Cell Line

Parental Cell line

HeLa

Genotype

Knockout

Gene Information

Gene Symbol

SLC38A2

Species

Human

Gene ID

54407

Swiss Prot

Q96QD8

Synonyms

ATA2; SAT2; SNAT2; PRO1068

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

Enables neutral amino acid:sodium symporter activity. Involved in several processes, including amino acid transport; cellular response to arsenite(3-); and positive regulation of RNA splicing. Located in cytoplasm and plasma membrane.

Product Information

Description

SLC38A2 Knockout cell line is engineered from HeLa cell line with Gene-Editing Technology.

Allele-1:79bp deletion in exon4

Allele-2:79bp deletion in exon4

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions

Dry ice

Amount

1~5x10⁶ cells/vial.

Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

Protocol

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO₂ condition.

1. Thaw the vial in 37°C water bath ,and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO₂.
7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

WT TTTTCCTCCAGCTA*****GTAAGGTACTCAT
Mut TTTTCCTCCAGCTA***Deletion***GTAAGGTACTCAT
Allele-1: 79bp deletion in exon4

WT TTTTCCTCCAGCTA*****GTAAGGTACTCAT
Mut TTTTCCTCCAGCTA***Deletion***GTAAGGTACTCAT
Allele-2: 79bp deletion in exon4

Genome sequence analysis of PCR products from parental (WT) and SLC38A2 knockout (KO) HeLa cells, using sanger sequencing.