

RBM15 Knockout HeLa Cell Line, Homozygous

Catalog No.: RM02687

Basic Information

Catalog No.

RM02687

Category

Cell Line

Parental Cell line

HeLa

Genotype

Knockout

Gene Information

Gene Symbol

RBM15

Species

Human

Gene ID

64783

Swiss Prot

Q96T37

Synonyms

OTT; OTT1; SPEN; RBM15

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

Members of the SPEN (Split-end) family of proteins, including RBM15, have repressor function in several signaling pathways and may bind to RNA through interaction with spliceosome components (Hiriart et al., 2005 [PubMed 16129689]).

Product Information

Description

RBM15 Knockout cell line is engineered from HeLa cell line with Gene-Editing Technology.

Allele-1:134bp deletion in exon1

Allele-2:134bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions

Dry ice

Amount

1~5x10⁶ cells/vial.

Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

Protocol

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO₂ condition.

1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO₂.
7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

WT CGACTTCCGCGGT*****TACCGGTGGGGGC
Mut CGACTTCCGCGGT***Deletion***TACCGGTGGGGGC
Allele-1: 134bp deletion in exon1

WT CGACTTCCGCGGT*****TACCGGTGGGGGC
Mut CGACTTCCGCGGT***Deletion***TACCGGTGGGGGC
Allele-2: 134bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and RBM15 knockout (KO) HeLa cells, using sanger sequencing.