ABclonal www.abclonal.com

RABEP1 Knockout 293T Cell Line, Homozygous

Catalog No.: RM02696

Basic Information

Catalog No.

RM02696

Category

Cell Line

Parental Cell line

293T

Genotype

Knockout

Background

Enables protein domain specific binding activity and protein homodimerization activity. Involved in vesicle-mediated transport. Located in endocytic vesicle and endosome. Part of protein-containing complex.

Gene Information

Gene Symbol

RABEP1

Species

Human

Gene ID

9135

Swiss Prot

Q15276

Synonyms

RAB5EP; RABPT5; RABEP1

Contact

2	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Product Information

Description

RABEP1 Knockout cell line is engineered from 293T cell line with Gene-Editing Technology. Allele-1:exon1 was deleted

Allele-2:exon1 was deleted

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions

Amount

Dry ice

1~5x10⁶ cells/vial.

Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

Protocol

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO_2 condition.

- 1. Thaw the vial in 37°C water bath ,and shake it to melt as soon as possible.
- 2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
- 3. Remove and discard the supernatant.
- 4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
- 5. Add 8-10mL of complete medium.
- 6. Incubate the culture at 37°C incubator with 5% CO₂.
- 7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

WT CCCGCTCCCCGAGG*************GCGGGCAGGCGGC
Mut CCCGCTCCCCGAGG***Deletion***GCGGGCAGGCGGC
Allele-1: exon1 was deleted

WT CCCGCTCCCCGAGG*************GCGGGCAGGCGGC
Mut CCCGCTCCCCGAGG***Deletion***GCGGGCAGGCGGC
Allele-2: exon1 was deleted

Genome sequence analysis of PCR products from parental (WT) and RABEP1 knockout (KO) 293T cells, using sanger sequencing.