Leader in Biomolecular Solutions for Life Science

www.abclonal.com

CHP1 Knockout 293T Cell Line, Homozygous

Catalog No.: RM02754

Basic Information

Catalog No.
RM02754

Category
Cell Line

Parental Cell line
293T

Genotype
Knockout

Background

This gene encodes a phosphoprotein that binds to the Na+/H+ exchanger NHE1. This protein serves as an essential cofactor which supports the physiological activity of NHE family members and may play a role in the mitogenic regulation of NHE1. The protein shares similarity with calcineurin B and calmodulin and it is also known to be an endogenous inhibitor of calcineurin activity.

Gene Information

Gene Symbol
CHP1

Species
Human

Gene ID
11261

Swiss Prot
Q99653

Synonyms
CHP; p22; p24; SPAX9; Sid470p; SLC9A1BP; CHP1

Product Information

Description
CHP1 Knockout cell line is engineered from 293T cell line with Gene-Editing Technology.
Allele-1:exon1 was deleted
Allele-2:exon1 was deleted
Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging
1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions
Dry ice

Amount
1~5x10^6 cells/vial.

Storage
Store at -20℃

Protocol
Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37℃ with 5% CO_2 condition.
1. Thaw the vial in 37℃ water bath ,and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37℃ incubator with 5% CO_2.
7. A subcultivation ratio of 1:2-1:4 is recommended.

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Sequencing data

```
WT    GGAGCTCCCGGCAC**********TAAGGTCTGGAGC
Mut   GGAGCTCCCGGCAC***Deletion***TAAGGTCTGGAGC
Allele-1:    exon1 was deleted

WT    AGGAGCTCCCGGCA**********TAAGGTCTGGAGC
Mut   AGGAGCTCCCGGCA***Deletion***TAAGGTCTGGAGC
Allele-2:    exon1 was deleted
```

Genome sequence analysis of PCR products from parental (WT) and CHP1 knockout (KO) 293T cells, using sanger sequencing.