

# PITX2 Knockout HeLa Cell Line, Homozygous

**Catalog No.:** RM02759

## Basic Information

**Catalog No.**

RM02759

**Category**

Cell Line

**Parental Cell line**

HeLa

**Genotype**

Knockout

## Gene Information

**Gene Symbol**

PITX2

**Species**

Human

**Gene ID**

5308

**Swiss Prot**

Q99697

**Synonyms**RS; RGS; ARP1; Brx1; IDG2; IGDS; IHG2;  
PTX2; RIEG; ASGD4; IGDS2; IRID2; Otlx2;  
RIEG1; Pituitary homeobox 2 (PITX2)

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

This gene encodes a member of the RIEG/PITX homeobox family, which is in the bicoid class of homeodomain proteins. The encoded protein acts as a transcription factor and regulates procollagen lysyl hydroxylase gene expression. This protein plays a role in the terminal differentiation of somatotroph and lactotroph cell phenotypes, is involved in the development of the eye, tooth and abdominal organs, and acts as a transcriptional regulator involved in basal and hormone-regulated activity of prolactin. Mutations in this gene are associated with Axenfeld-Rieger syndrome, iridogoniodysgenesis syndrome, and sporadic cases of Peters anomaly. A similar protein in other vertebrates is involved in the determination of left-right asymmetry during development. Alternatively spliced transcript variants encoding distinct isoforms have been described.

## Product Information

**Description**

PITX2 Knockout cell line is engineered from HeLa cell line with Gene-Editing Technology.

Allele-1:139bp deletion in exon2

Allele-2:154bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

**Packaging**

1 vial parental cell line and 1 vial knockout cell line

**Shipping Conditions**

Dry ice

**Amount**1~5x10<sup>6</sup> cells/vial.**Storage**

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

**Protocol**

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO<sub>2</sub> condition.

1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO<sub>2</sub>.
7. A subcultivation ratio of 1:2-1:4 is recommended.

## Sequencing data

---

WT TCTAAGAAGAAGCG\*\*\*\*\*ATCTCAGAGCTGG  
Mut TCTAAGAAGAAGCG\*\*\*Deletion\*\*\*ATCTCAGAGCTGG  
Allele-1: 139bp deletion in exon2

WT GCGCGCGAGGACCC\*\*\*\*\*AATGTGGAAGGCAG  
Mut GCGCGCGAGGACCC\*\*\*Deletion\*\*\*AATGTGGAAGGCAG  
Allele-2: 154bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and PITX2 knockout (KO) HeLa cells, using sanger sequencing.