

CD1A Knockout 293T Cell Lysate, Homozygous

Catalog No.: RM02782

Basic Information

Catalog No.

RM02782

Category

Cell Lysate

Parental Cell line

293T

Genotype

Knockout

Gene Information

Gene Symbol

CD1A

Species

Human

Gene ID

909


Swiss Prot

P06126

Synonyms

R4; T6; CD1; FCB6; HTA1; CD1A

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

This gene encodes a member of the CD1 family of transmembrane glycoproteins, which are structurally related to the major histocompatibility complex (MHC) proteins and form heterodimers with beta-2-microglobulin. The CD1 proteins mediate the presentation of primarily lipid and glycolipid antigens of self or microbial origin to T cells. The human genome contains five CD1 family genes organized in a cluster on chromosome 1. The CD1 family members are thought to differ in their cellular localization and specificity for particular lipid ligands. The protein encoded by this gene localizes to the plasma membrane and to recycling vesicles of the early endocytic system. Alternative splicing results in multiple transcript variants.

Product Information

Description

CD1A Knockout cell line is engineered from 293T cell line with Gene-Editing Technology.

Allele-1:112bp deletion in exon2

Allele-2:112bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50µL, 2µg/µL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT ATTGCAGACTCAT*****GGTCATTGAGGGA
Mut ATTGCAGACTCAT***Deletion***GGTCATTGAGGGA
Allele-1: 112bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and CD1A knockout (KO) 293T cells, using sanger sequencing.

WT ATTGCAGACTCAT*****GGTCATTGAGGGA
Mut ATTGCAGACTCAT***Deletion***GGTCATTGAGGGA
Allele-2: 112bp deletion in exon2