

PAX5 Knockdown A549 Cell Lysate, Heterozygous

Catalog No.: RM02785

Basic Information

Catalog No.

RM02785

Category

Cell Lysate

Parental Cell line

A549

Genotype

Knockdown

Gene Information

Gene Symbol

PAX5

Species

Human

Gene ID

5079

Swiss Prot

Q02548

Synonyms

ALL3; BSAP; PAX-5; PAX5

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

This gene encodes a member of the paired box (PAX) family of transcription factors. The central feature of this gene family is a novel, highly conserved DNA-binding motif, known as the paired box. Paired box transcription factors are important regulators in early development, and alterations in the expression of their genes are thought to contribute to neoplastic transformation. This gene encodes the B-cell lineage specific activator protein that is expressed at early, but not late stages of B-cell differentiation. Its expression has also been detected in developing CNS and testis and so the encoded protein may also play a role in neural development and spermatogenesis. This gene is located at 9p13, which is involved in t(9;14)(p13;q32) translocations recurring in small lymphocytic lymphomas of the plasmacytoid subtype, and in derived large-cell lymphomas. This translocation brings the potent E-mu enhancer of the IgH gene into close proximity of the PAX5 promoter, suggesting that the deregulation of transcription of this gene contributes to the pathogenesis of these lymphomas. Alternative splicing results in multiple transcript variants encoding different isoforms.

Product Information

Description

PAX5 Knockdown A549 cell line is engineered from A549 cell line with Gene-Editing Technology.

Allele-1: exon4 was deleted

Allele-2: WT

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50µL, 2µg/µL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT TCTCTTCCTTCC*****GCTGAACCTGGTAA
Mut TCTCTTCCTTCC***Deletion***GCTGAACCTGGTAA
Allele-1: exon4 was deleted

WT TCTCTTCCTTCC*****GCTGAACCTGGTAA
Mut TCTCTTCCTTCC*****GCTGAACCTGGTAA
Allele-2: WT

Genome sequence analysis of PCR products from parental (WT) and PAX5 knockdown (KD) A549 cells, using sanger sequencing.