Rabbit anti-Human IL-10 mAb (CAP)



Catalog No.: RM17643

Basic Information

Catalog No.

RM17643

Catagory

Elisa Antibody Kit

Application

Multiplex,CLIA

Product Information

Ig Type

Rabbit IgG

Purification

Affinity purification

Endotoxin Level

Storage

Store at -20 $^{\circ}\text{C}.$

Avoid repeated freeze-thaw cycles.

Formulation

Supplied as a 0.2µm filtered solution in PBS with 0.05% Proclin300, PH 7.4.

Contact

•	order@abclonal.com
<u>a</u>	support@abclonal.com
•	www.abclonal.com

Background

The protein encoded by this gene is a cytokine produced primarily by monocytes and to a lesser extent by lymphocytes. This cytokine has pleiotropic effects in immunoregulation and inflammation. It down-regulates the expression of Th1 cytokines, MHC class II Ags, and costimulatory molecules on macrophages. It also enhances B cell survival, proliferation, and antibody production. This cytokine can block NF-kappa B activity, and is involved in the regulation of the JAK-STAT signaling pathway. Knockout studies in mice suggested the function of this cytokine as an essential immunoregulator in the intestinal tract. Mutations in this gene are associated with an increased susceptibility to HIV-1 infection and rheumatoid arthritis.

Immunogen Information

Immunogen

Recombinant Human IL-10 Protein Ser19-Asn178 Accession #P22301

Cross-Reactivity


No cross-reactivity in multiplex assay with recombinant rIL-10, rIL-10

Assay Applications


Human IL-10 Sandwich Immunoassay

		Recommended Concentration	Sample
	Capture	1-4ug/mL	Rabbit anti-Human IL-10 mAb (Cat. No. RM17643)
CLIA	Detection	0.5-2ug/mL	Rabbit anti-Human IL-10 mAb (Cat. No. RM17644)
	Standard	2.5-1000pg/mL	Recombinant Human IL-10 Protein

	Capture	1-4ug/mL	Rabbit anti-Human IL-10 mAb (Cat. No. RM17643)
Multiplex	Detection	0.5-2ug/mL	Rabbit anti-Human IL-10 mAb (Cat. No. RM17644)
	Standard	1.4-1000pg/mL	Recombinant Human IL-10 Protein



This standard curve is only for demonstration purposes. A standard curve should be generated for each assay.



This standard curve is only for demonstration purposes. A standard curve should be generated for each assay.