
Leader in Biomolecular Solutions for Life Science

# Rabbit anti-Human IL-12 p70 mAb



## **Basic Information**

Catalog No. RM17739

**Catagory** Elisa Antibody Kit

Application ELISA,Multiplex,CLIA

# **Product Information**

**Ig Type** Rabbit IgG

**Purification** Affinity purification

#### **Endotoxin Level**

Storage Store at -20°C.

Avoid repeated freeze-thaw cycles.

#### Formulation

Supplied as a  $0.2 \mu m$  filtered solution in PBS with 0.05% Proclin300, PH 7.4.

### Contact

| - | order@abclonal.com   |
|---|----------------------|
| 6 | support@abclonal.com |
| Ð | www.abclonal.com     |

#### Background

IL-12 is a heterodimeric cytokine composed of p40 and p35 subunits and is considered a largely pro-inflammatory cytokine. It is naturally produced by dendritic cells, macrophages, neutrophils, and human B-lymphoblastoid cells. IL-12?is crucial for the recruitment and effector functions of CD8+?T and NK cells.

### **Immunogen Information**

Immunogen Recombinant IL-12 p70 Protein

# **Cross-Reactivity**

No cross-reactivity in ELISA assay with recombinant hG-CSF[]hIL-6[]rIL-6


# **Assay Applications**

Human IL-12 p70 ELISA/CLIA


|       |           | Recommended<br>Concentration | Sample   |
|-------|-----------|------------------------------|--|
| ELISA | Capture   | 0.5-2ug/mL                   | Rabbit anti-Human IL-12 p70 mAb(Cat. No.RM17739)         |
|       | Detection | 0.02-0.08ug/mL               | Rabbit anti-Human IL-12 p70 mAb(Cat. No.RM17740]         |
|       | Standard  | 31.25-2000pg/mL              | Recombinant Human IL-12 p70<br>Protein(Cat. No.RP01232[] |




| CLIA | Capture   | 1-4ug/mL    | Rabbit anti-Human IL-12 p70 mAb(Cat.<br>No.RM17746) |
|------|-----------|-------------|---|
|      | Detection | 0.5-2ug/mL  | Rabbit anti-Human IL-12 p70 mAb(Cat.<br>No.RM17739) |
|      | Standard  | 0.1-10ng/mL | Recombinant Human IL-12 p70 Protein                 |




This standard curve is only for demonstration purposes. A standard curve should be generated for eachassay.




No significant cross-reactivity or interference was observed.



No significant cross-reactivity or interference was observed.



Placed at  $37^{\circ}$ C for 3 days, the stability of the standard curve all conform to CV <10%.



This standard curve is only for demonstration purposes. A standard curve should be generated for eachassay.