Leader in Biomolecular Solutions for Life Science

www.abclonal.com

Rabbit anti-Human sST2 mAb

Catalog No.: RM17779

Basic Information

Catalog No.
RM17779

Catagory
Elisa Antibody Kit

Application
ELISA

Product Information

Ig Type
Rabbit IgG

Purification
Affinity purification

Endotoxin Level

Storage
Store at -20℃.
Avoid repeated freeze-thaw cycles.

Formulation
Supplied as a 0.2μm filtered solution in PBS with 0.05% Proclin300, PH 7.4.

Contact

order@abclonal.com

support@abclonal.com

www.abclonal.com

Background

Receptor for somatostatin-14 and -28. This receptor is coupled via pertussis toxin sensitive G proteins to inhibition of adenylyl cyclase. In addition it stimulates phosphotyrosine phosphatase and PLC via pertussis toxin insensitive as well as sensitive G proteins. Inhibits calcium entry by suppressing voltage-dependent calcium channels. Acts as the functionally dominant somatostatin receptor in pancreatic alpha- and beta-cells where it mediates the inhibitory effect of somatostatin-14 on hormone secretion. Inhibits cell growth through enhancement of MAPK1 and MAPK2 phosphorylation and subsequent up-regulation of CDKN1B. Stimulates neuronal migration and axon outgrowth and may participate in neuron development and maturation during brain development. Mediates negative regulation of insulin receptor signaling through PTPN6. Inactivates SSTR3 receptor function following heterodimerization

Immunogen Information

Immunogen
Recombinant Human sST2 Protein

Cross-Reactivity

No significant cross-reactivity or interference between sST2 and analogues was observed.

Assay Applications

Human sST2 Sandwich Immunoassay

		Recommended Concentration	Sample
ELISA	Capture	1-4ug/mL	Rabbit anti-Human sST2 mAb(Cat. No.RM17778)
	Detection	0.015-0.06ug/mL	Rabbit anti-Human sST2 mAb(Cat. No.RM17779)
	Standard	7.8-500pg/mL	Recombinant Human sST2 Protein

Validation Data



This standard curve is only for demonstration purposes. A standard curve should be generated for eachassay.

Fromthe comparison of the standard curve, it is found that ABclonal has highsensitivity.



No significant cross-reactivity or interference was observed.



Placed at 37°C for 3 days, the stability of the standard curve all conformto CV <10%.