Rabbit anti-Rat IL-10 mAb



Catalog No.: RM17825

Basic Information

Catalog No.

RM17825

Catagory

Elisa Antibody Kit

Application

ELISA

Product Information

Ig Type

Rabbit IgG

Purification

Affinity purification

Endotoxin Level

Storage

Store at -20°C.

Avoid repeated freeze-thaw cycles.

Formulation

Supplied as a 0.2um filtered solution in PBS with 0.05%ProClin 300,PH 7.4.

Contact

:		order@abclonal.com
2		support@abclonal.com
•	Ī	www.abclonal.com

Background

Enables interleukin-10 receptor binding activity. Involved in several processes, including cellular response to estradiol stimulus; liver regeneration; and negative regulation of nitric oxide biosynthetic process. Located in extracellular space. Used to study several diseases, including lung disease (multiple); perinatal necrotizing enterocolitis; sciatic neuropathy; toxic encephalopathy; and transient cerebral ischemia. Biomarker of several diseases, including artery disease (multiple); decubitus ulcer; lung disease (multiple); obstructive sleep apnea; and postpartum depression. Human ortholog(s) of this gene implicated in several diseases, including artery disease (multiple); autoimmune disease (multiple); eye disease (multiple); gastrointestinal system cancer (multiple); and hematologic cancer (multiple). Orthologous to human IL10 (interleukin 10).

Immunogen Information

Immunogen

Recombinant Rat IL-10


Cross-Reactivity

No cross-reactivity in ELISA assay withrecombinant Human IL-10,Mouse IL-10,Rat IFNG,Rat IL-1B,Rat IL-2,Rat IL-4,Rat TNFA


Assay Applications

Rat IL-10 Sandwich ELISA


		Recommended Concentration	Sample
ELISA	Capture	1-4ug/mL	Rabbit anti-Rat IL-10 mAb(Cat. No.RM17825)
	Detection	0.0625-0.25ug/mL	Rabbit anti-Rat IL-10 mAb(Cat. No.RM17826)
	Standard	7.81-500pg/mL	Recombinant Rat IL-10 Protein



This standard curve is only for demonstration purposes. A standard curve should be generated for each assay.



Fromthe comparison of the standard curve, it is found that ABclonal has highsensitivity.



Placed at 37°C for 3 days, the stability of the standard curve all conform to CV <10%.