# Rabbit anti-Human IL-2R mAb



Catalog No.: RM17893

# **Basic Information**

### Catalog No.

RM17893

#### Catagory

Elisa Antibody Kit

#### **Application**

CLIA

### **Product Information**

# Ig Type

Rabbit IgG

# **Purification**

Affinity purification

### **Endotoxin Level**

#### Storage

Store at -20°C.

Avoid repeated freeze-thaw cycles.

# **Formulation**

Supplied as a 0.2µm filtered solution in PBS with 0.05% Proclin300, PH 7.4.

### **Contact**

•	order@abclonal.com
<u>a</u>	support@abclonal.com
•	www.abclonal.com

# **Background**

The interleukin 2 (IL2) receptor alpha (IL2RA) and beta (IL2RB) chains, together with the common gamma chain (IL2RG), constitute the high-affinity IL2 receptor. Homodimeric alpha chains (IL2RA) result in low-affinity receptor, while homodimeric beta (IL2RB) chains produce a medium-affinity receptor. Normally an integral-membrane protein, soluble IL2RA has been isolated and determined to result from extracellular proteolyisis. Alternately-spliced IL2RA mRNAs have been isolated, but the significance of each is presently unknown. Mutations in this gene are associated with interleukin 2 receptor alpha deficiency. Patients with severe Coronavirus Disease 2019 (COVID-19), the disease caused by the novel severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), have significantly elevated levels of IL2R in their plasma. Similarly, serum IL-2R levels are found to be elevated in patients with different types of carcinomas. Certain IL2RA and IL2RB gene polymorphisms have been associated with lung cancer risk.

# **Immunogen Information**

### **Immunogen**


Recombinant Human IL-2R alpha Protein

# **Cross-Reactivity**

# **Assay Applications**

Human IL-2R Sandwich Immunoassay

		Recommended Concentration	Sample
CLIA	Capture	1-4ug/mL	Rabbit anti-Human IL-2R mAb(Cat. No.RM17892)
	Detection	0.5-2ug/mL	Rabbit anti-Human IL-2R mAb(Cat. No.RM17893)
	Standard	5-2000pg/mL	Recombinant Human IL-2R Protein



This standard curve is only for demonstration purposes. A standard curve should be generated for each assay.