

Rabbit anti-Mouse IL-10 mAb

Catalog No.: RM17935

Basic Information

Catalog No.
RM17935

Catagory
Elisa Antibody Kit

Application
ELISA, Multiplex

Product Information

Ig Type
Rabbit IgG

Purification
Affinity purification

Endotoxin Level
/

Storage
Store at -20°C.
Avoid repeated freeze-thaw cycles.

Formulation
Supplied as a 0.2 µm filtered solution in PBS with 0.05% Proclin 300, pH 7.4.

Contact

 | order@abclonal.com

 | support@abclonal.com

 | www.abclonal.com

Background

This gene encodes an anti-inflammatory cytokine that is a member of the class-2 cytokine family. The encoded protein is secreted by cells of both the innate and adaptive immune systems and is crucial for limiting the immune response to a broad range of pathogens. It also has been shown to suppress autoimmune responses. This protein mediates its immunosuppressive signal through a specific interleukin 10 receptor complex. Aberrant functioning of this gene is associated with numerous immune disorders including graft-versus-host disease, and increased susceptibility to HIV-1 infection and rheumatoid arthritis.

Immunogen Information

Immunogen
Recombinant Mouse IL-10

Cross-Reactivity

No cross-reactivity in immunoassay with recombinant Mouse IFN-γ, Mouse IFN-β, Mouse IL-20, Mouse IL-22, Mouse IL-24 and Human IL-10. Cross-reactivity was found with Rat IL-10.


Assay Applications

Mouse IL-10 Sandwich Immunoassay


		Recommended Concentration	Sample
ELISA	Capture	0.5-2 µg/mL	Rabbit anti-Mouse IL-10 mAb (Cat. No. RM17935)
	Detection	0.015-0.06 µg/mL	Rabbit anti-Mouse IL-10 mAb (Cat. No. RM17936)
	Standard	31.25-2000 pg/mL	Recombinant Mouse IL-10 Protein (Cat. No. RP01465)

Multiplex	Capture	0.1-5 µg/mL	Rabbit anti-Mouse IL-10 mAb (Cat. No. RM17935)
	Detection	0.5-2 µg/mL	Rabbit anti-Mouse IL-10 mAb (Cat. No. RMK0423)
	Standard	6.86-5000 pg/mL	Recombinant Mouse IL-10 Protein (Cat. No. RP01465)


Validation Data



This standard curve is only for demonstration purposes. A standard curve should be generated for each assay.



No significant cross-reactivity or interference was observed.



Placed at 37°C for 3 days, the stability of the standard curve all conform to CV <10%.