Rabbit anti-Mouse IL-10 mAb



Catalog No.: RM17935

Basic Information

Catalog No.

RM17935

Catagory

Elisa Antibody Kit

Application

ELISA, Multiplex

Product Information

Ig Type

Rabbit IgG

Purification

Affinity purification

Endotoxin Level

1

Storage

Store at -20°C.

Avoid repeated freeze-thaw cycles.

Formulation

Supplied as a 0.2 µm filtered solution in PBS with 0.05% Proclin 300, pH 7.4.

Contact

②	www.abclonal.com
a	support@abclonal.com
<u>•</u>	order@abclonal.com

Background

This gene encodes an anti-inflammatory cytokine that is a member of the class-2 cytokine family. The encoded protein is secreted by cells of both the innate and adaptive immune systems and is crucial for limiting the immune response to a broad range of pathogens. It also has been shown to suppress autoimmune responses. This protein mediates it's immunosuppressive signal through a specific interleukin 10 receptor complex. Aberrant functioning of this gene is associated with numerous immune disorders including graft-versus-host disease, and increased susceptibility to HIV-1 infection and rheumatoid arthritis.

Immunogen Information

Immunogen

Recombinant Mouse IL-10

Cross-Reactivity


No cross-reactivity in immunoassay with recombinant Mouse IFN- γ , Mouse IFN- β , Mouse IL-20, Mouse IL-22, Mouse IL-24 and Human IL-10. Cross-reactivity was found with Rat IL-10.

Assay Applications


Mouse IL-10 Sandwich Immunoassay

		Recommended Concentration	Sample
ELISA	Capture	0.5-2 μg/mL	Rabbit anti-Mouse IL-10 mAb (Cat. No. RM17935)
	Detection	0.015-0.06 μg/mL	Rabbit anti-Mouse IL-10 mAb (Cat. No. RM17936)
	Standard	31.25-2000 pg/mL	Recombinant Mouse IL-10 Protein (Cat. No. RP01465)


Multiplex	Capture	0.1-5 μg/mL	Rabbit anti-Mouse IL-10 mAb (Cat. No. RM17935)
	Detection	0.5-2 μg/mL	Rabbit anti-Mouse IL-10 mAb (Cat. No. RMK0423)
	Standard	6.86-5000 pg/mL	Recombinant Mouse IL-10 Protein (Cat. No. RP01465)



This standard curve is only for demonstration purposes. A standard curve should be generated for each assay.



No significant cross-reactivity or interference was observed.



Placed at 37°C for 3 days, the stability of the standard curve all conform to CV <10%.