Rabbit anti-Human IL-12 p70 mAb(DET[]



Catalog No.: RM17972

Basic Information

Catalog No.

RM17972

Catagory

Elisa Antibody Kit

Application

multiplex assay

Product Information

Ig Type

Rabbit IgG

Purification

Affinity purification

Endotoxin Level

Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free.

Avoid repeated freeze-thaw cycles.

Formulation

Supplied as a 0.2um filtered solution in PBS,PH 7.4 containing Human IL-12p70 Antibody.

Contact

| • | order@abclonal.com |
|----------|----------------------|
| <u>a</u> | support@abclonal.com |
| • | www.abclonal.com |

Background

Interleukin 12 (IL-12), also known as natural killer cell stimulatory factor (NKSF) or cytotoxic lymphocyte maturation factor (CLMF), is a heterodimeric pleiotropic cytokine made up of a 40 kDa (p40) subunit and a 35 kDa (p35) subunit. The IL-12 p40 subunit is shared by IL-23, another heterodimeric cytokine that has biological activities similar to, as well as distinct from, IL-12. IL-12 is produced by macrophages and B cells and has been shown to have multiple effects on T cells and natural killer (NK) cells. While mouse IL-12 is active on both human and mouse cells, human IL-12 is not active on mouse cells.

Immunogen Information

Immunogen


Recombinant Human IL-12 p70 Protein IL12A(Arg23-Ser219)/IL12B (Ile23-Ser328) Accession #P29460 P29459

Cross-Reactivity

Assay Applications

Human IL-12p70 multiplex assay

| | Recommended Concentration | Sample |
|------------------------|------------------------------|---|
| Multiplex Capture | 3-20ug/mL | Rabbit anti-Human IL-12p70 mAb(CAP)(Cat. No.RM17971 |
| Multiplex Detection | 0.017-2ug/mL | Rabbit anti-Human IL-12p70 mAb(DET)(Cat. No.RM17972) |
| Standard | 13.72-10000pg/mL | Recombinant Human IL-12p70 Protein (Cat. No. RP01232) |



This standard curve is only for demonstration purposes. A standard curve should be generated for each assay.