# ABclonal www.abclonal.com

# GDF3 Knockout 293T Cell Lysate, Homozygous

Catalog No.: RM50006

#### **Basic Information**

#### Catalog No.

RM50006

#### Category

Cell Lysate

#### **Parental Cell line**

293T

#### Genotype

Knockout

## **Background**

This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked homodimer. This protein plays a role ocular and skeletal development. Mutations in this gene are associated with microphthalmia, coloboma, and skeletal abnormalities in human patients.

#### **Gene Information**

#### **Gene Symbol**

GDF3

#### **Species**

Human

# Gene ID

9573

#### **Swiss Prot**

Q9NR23

#### **Synonyms**

KFS3; MCOP7; MCOPCB6; GDF3

#### **Contact**

2	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

### **Product Information**

#### Description

GDF3 Knockout cell line is engineered from 293T cell line with Gene-Editing Technology. Allele-1:82bp deletion in exon1

Allele-2:82bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

## **Packaging**

1 vial parental cell Lysate and 1 vial knockout cell Lysate

# **Shipping Conditions**

**Amount** 

4°C

50μL, 2μg/μL.

#### Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

## Protocol

To be used as WB control. Lysate is supplied in  $1\times$  SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

# Sequencing data

WT ATTTCTGGGCTTAG\*\*\*\*\*\*\*\*\*\*\*\*\*\*CACTGGGGTCTCCC
Mut ATTTCTGGGCTTAG\*\*\*Deletion\*\*\*CACTGGGGTCTCCC
Allele-1: 82bp deletion in exon1

WT ATTTCTGGGCTTAG\*\*\*\*\*\*\*\*\*\*\*\*CACTGGGGTCTCCC
Mut ATTTCTGGGCTTAG\*\*\*Deletion\*\*\*CACTGGGGTCTCCC
Allele-2: 82bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and GDF3 knockout (KO) 293T cells, using sanger sequencing.