# ABclonal www.abclonal.com

## ABL2 Knockout 293T Cell Lysate, Homozygous

Catalog No.: RM50007

#### **Basic Information**

#### Catalog No.

RM50007

#### Category

Cell Lysate

#### **Parental Cell line**

293T

#### Genotype

Knockout

#### **Background**

This gene encodes a member of the Abelson family of nonreceptor tyrosine protein kinases. The protein is highly similar to the c-abl oncogene 1 protein, including the tyrosine kinase, SH2 and SH3 domains, and it plays a role in cytoskeletal rearrangements through its C-terminal F-actin- and microtubule-binding sequences. This gene is expressed in both normal and tumor cells, and is involved in translocation with the ets variant 6 gene in leukemia. Multiple alternatively spliced transcript variants encoding different protein isoforms have been found for this gene.

#### **Gene Information**

#### **Gene Symbol**

ABL2

#### **Species**

Human

#### Gene ID

27

#### **Swiss Prot**

P42684

#### Synonyms

ARG; ABLL; ABL2

#### **Contact**

<u>a</u>		400-999-6126
$\sim$		cn.market@abclonal.com.cn
•	Τ	www.abclonal.com.cn

#### **Product Information**

#### Description

ABL2 Knockout cell line is engineered from 293T cell line with Gene-Editing Technology. Allele-1:94bp deletion in exon3

Allele-2:94bp deletion in exon3

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

#### **Packaging**

1 vial parental cell Lysate and 1 vial knockout cell Lysate

# Shipping Conditions

**Amount** 50μL, 2μg/μL.

#### Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

#### **Protocol**

To be used as WB control. Lysate is supplied in  $1\times$  SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

### Sequencing data

WT ACCATGGACCTGTG\*\*\*\*\*\*\*\*\*\*\*\*\*CGCTCAGGTACGA
Mut ACCATGGACCTGTG\*\*\*Deletion\*\*\*CGCTCAGGTACGA
Allele-1: 94bp deletion in exon3

WT ACCATGGACCTGTG\*\*\*\*\*\*\*\*\*\*\*CGCTCAGGTACGA
Mut ACCATGGACCTGTG\*\*\*Deletion\*\*\*CGCTCAGGTACGA

Allele-1: 94bp deletion in exon3

Genome sequence analysis of PCR products from parental (WT) and ABL2 knockout (KO) 293T cells, using sanger sequencing.