

CD320 Knockout 293T Cell Lysate, Homozygous

Catalog No.: RM50010

Basic Information

Catalog No.

RM50010

Category

Cell Lysate

Parental Cell line

293T

Genotype

Knockout

Gene Information

Gene Symbol

CD320

Species

Human

Gene ID

51293

Swiss Prot

Q9NPF0


Synonyms

8D6; 8D6A; TCBLR; TCN2R; TCII-R;
sCD320; CD320

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

This gene encodes the transcobalamin receptor that is expressed at the cell surface. It mediates the cellular uptake of transcobalamin bound cobalamin (vitamin B12), and is involved in B-cell proliferation and immunoglobulin secretion. Mutations in this gene are associated with methylmalonic aciduria. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Product Information

Description

CD320 Knockout cell line is engineered from 293T cell line with Gene-Editing Technology.

Allele-1:64bp deletion in exon1

Allele-2:64bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50µL, 2µg/µL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT GGCGTTGGATG*****ACTAGCCTGGAG
Mut GGCGTTGGATG***Deletion***ACTAGCCTGGAG
Allele-1: 64bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and CD320 knockout (KO) 293T cells, using sanger sequencing.

WT GGCGTTGGATG*****ACTAGCCTGGAG
Mut GGCGTTGGATG***Deletion***ACTAGCCTGGAG
Allele-2: 64bp deletion in exon1