

NRG3 Knockout 293T Cell Lysate, Homozygous

Catalog No.: RM50012

Basic Information

Catalog No.

RM50012

Category

Cell Lysate

Parental Cell line

293T

Genotype

Knockout

Background

This gene is a member of the neuregulin gene family. This gene family encodes ligands for the transmembrane tyrosine kinase receptors ERBB3 and ERBB4 - members of the epidermal growth factor receptor family. Ligand binding activates intracellular signaling cascades and the induction of cellular responses including proliferation, migration, differentiation, and survival or apoptosis. This gene encodes neuregulin 3 (NRG3). NRG3 has been shown to activate the tyrosine phosphorylation of its cognate receptor, ERBB4, and is thought to influence neuroblast proliferation, migration and differentiation by signalling through ERBB4. NRG3 also promotes mammary differentiation during embryogenesis. Linkage studies have implicated this gene as a susceptibility locus for schizophrenia and schizoaffective disorder. Alternative splicing results in multiple transcript variants encoding distinct isoforms. Additional transcript variants have been described but their biological validity has not been verified.

Gene Information

Gene Symbol

NRG3

Species

Human

Gene ID

10718

Swiss Prot

P56975

Synonyms

HRG3; pro-NRG3; Neuregulin-3 (NRG3)

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Description

NRG3 Knockout cell line is engineered from 293T cell line with Gene-Editing Technology.

Allele-1:82bp deletion in exon3

Allele-2:83bp deletion in exon3

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50µL, 2µg/µL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT ATATTCCACAGAG*****TGACCGGATCCCAT
Mut ATATTCCACAGAG***Deletion***TGACCGGATCCCAT
Allele-1: 82bp deletion in exon3

Genome sequence analysis of PCR products from parental (WT) and NRG3 knockout (KO) 293T cells, using sanger sequencing.

WT CATATTCCACAGA*****TGACCGGATCCCAT
Mut CATATTCCACAGA***Deletion***TGACCGGATCCCAT
Allele-2: 83bp deletion in exon3