# ABclonal www.abclonal.com

## **HOPX Knockout 293T Cell Lysate, Homozygous**

Catalog No.: RM50013

#### **Basic Information**

#### Catalog No.

RM50013

#### Category

Cell Lysate

#### **Parental Cell line**

293T

#### Genotype

Knockout

#### **Background**

The protein encoded by this gene is a homeodomain protein that lacks certain conserved residues required for DNA binding. It was reported that choriocarcinoma cell lines and tissues failed to express this gene, which suggested the possible involvement of this gene in malignant conversion of placental trophoblasts. Studies in mice suggest that this protein may interact with serum response factor (SRF) and modulate SRF-dependent cardiac-specific gene expression and cardiac development. Multiple alternatively spliced transcript variants have been identified for this gene.

#### **Gene Information**

#### **Gene Symbol**

**HOPX** 

#### **Species**

Human

#### Gene ID

84525

#### **Swiss Prot**

Q9BPY8

#### **Synonyms**

HOD; HOP; OB1; LAGY; TOTO; CAMEO; NECC1; SMAP31; HOPX

#### **Contact**

6	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

#### **Product Information**

#### Description

HOPX Knockout cell line is engineered from 293T cell line with Gene-Editing Technology. Allele-1:95bp deletion in exon1

Allele-2:95bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

#### **Packaging**

1 vial parental cell Lysate and 1 vial knockout cell Lysate

# Shipping ConditionsAmount $4^{\circ}C$ $50\mu L$ , $2\mu g/\mu L$ .

#### Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

#### **Protocol**

To be used as WB control. Lysate is supplied in  $1\times$  SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

### Sequencing data

WT TCGGCGGAGACCGC\*\*\*\*\*\*\*\*\*\*\*\*\*CCGAGGCAGGCCT
Mut TCGGCGGAGACCGC\*\*\*Deletion\*\*\*CCGAGGCAGGCCT
Allele-1: 95bp deletion in exon1

WT TCGGCGGAGACCGC\*\*\*\*\*\*\*\*\*\*\*\*CCGAGGCAGGCCT
Mut TCGGCGGAGACCGC\*\*\*Deletion\*\*\*CCGAGGCAGGCCT

Allele-2: 95bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and HOPX knockout (KO) 293T cells, using sanger sequencing.