

IL1RN Knockout 293T Cell Lysate, Homozygous

Catalog No.: RM50016

Basic Information

Catalog No.

RM50016

Category

Cell Lysate

Parental Cell line

293T

Genotype

Knockout

Gene Information

Gene Symbol

IL1RN

Species

Human

Gene ID


3557

Swiss Prot

P18510

SynonymsDIRA; IRAP; IL1F3; IL1RA; MVCD4; IL-1RN;
IL-1ra; IL-1ra3; ICIL-1RA

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

The protein encoded by this gene is a member of the interleukin 1 cytokine family. This protein inhibits the activities of interleukin 1, alpha (IL1A) and interleukin 1, beta (IL1B), and modulates a variety of interleukin 1 related immune and inflammatory responses, particularly in the acute phase of infection and inflammation. This gene and five other closely related cytokine genes form a gene cluster spanning approximately 400 kb on chromosome 2. A polymorphism of this gene is reported to be associated with increased risk of osteoporotic fractures and gastric cancer. Several alternatively spliced transcript variants encoding distinct isoforms have been reported.

Product Information

Description

IL1RN Knockout cell line is engineered from 293T cell line with Gene-Editing Technology.

Allele-1:80bp deletion in exon1

Allele-2:89bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50µL, 2µg/µL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT AACTCTGGGCCG*****CTCTGGGAGAAAAT
Mut AACTCTGGGCCG***Deletion***CTCTGGGAGAAAAT
Allele-1: 80bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and IL1RN knockout (KO) 293T cells, using sanger sequencing.

WT AACTCTGGGCCG*****AAAATCCAGCAAGA
Mut AACTCTGGGCCG***Deletion***AAAATCCAGCAAGA
Allele-2: 89bp deletion in exon1