Leader in Biomolecular Solutions for Life Science

www.abclonal.com

DCTN4 Knockdown HeLa Cell Line, Heterozygous

Catalog No.: RM50047

Basic Information

Catalog No.
RM50047

Category
Cell Line

Parental Cell line
HeLa

Genotype
Knockdown

Gene Information

Gene Symbol
DCTN4

Species
Human

Gene ID
51164

Swiss Prot
Q9UJW0

Synonyms
P62; DYN4

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Background

Enables protein N-terminus binding activity. Located in centrosome. [provided by Alliance of Genome Resources, Apr 2022]

Product Information

Description
DCTN4 Knockdown HeLa cell line is engineered from HeLa cell line with Gene-Editing Technology.
Allele-1:6bp deletion in exon3
Allele-2:166bp deletion in exon3
Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging
1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions
Dry ice

Amount
1~5x10^6 cells/vial.

Storage
Stored in liquid nitrogen for a long time less than -130℃. Minimizing freeze-thaw cycles.

Protocol
Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37℃ with 5% CO_2 condition.
1. Thaw the vial in 37℃ water bath ,and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37℃ incubator with 5% CO_2.
7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data



Genome sequence analysis of PCR products from parental (WT) and DCTN4 knockdown (KD) HeLa cells, using sanger sequencing.