

SFRP2 Knockout HeLa Cell Line, Homozygous

Catalog No.: RM50059

Basic Information

Catalog No.

RM50059

Category

Cell Line

Parental Cell line

HeLa

Genotype

Knockout

Gene Information

Gene Symbol

SFRP2

Species

Human

Gene ID

6423

Swiss Prot

Q96HF1

Synonyms

FRP-2; SARP1; SDF-5; SFRP2

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

This gene encodes a member of the SFRP family that contains a cysteine-rich domain homologous to the putative Wnt-binding site of Frizzled proteins. SFRPs act as soluble modulators of Wnt signaling. Methylation of this gene is a potential marker for the presence of colorectal cancer.

Product Information

Description

SFRP2 Knockout cell line is engineered from HeLa cell line with Gene-Editing Technology.

Allele-1:97bp deletion in exon1

Allele-2:34bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions

Dry ice

Amount

1~5x10⁶ cells/vial.

Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

Protocol

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO₂ condition.

1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO₂.
7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

WT CCTTCGGCTTCCCC*****ACTCCCTGCTGGGC
Mut CCTTCGGCTTCCCC***Deletion***ACTCCCTGCTGGGC
Allele-1: 97bp deletion in exon1

WT GGCTTCCCCTGGCC*****AAGACCTTTGCAT
Mut GGCTTCCCCTGGCC***Deletion***AAGACCTTTGCAT
Allele-2: 34bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and SFRP2 knockout (KO) HeLa cells, using sanger sequencing.