Leader in Biomolecular Solutions for Life Science

www.abclonal.com

ZIP14 Knockout HeLa Cell Line, Homozygous

Catalog No.: RM50071

Basic Information

Catalog No.
RM50071

Category
Cell Line

Parental Cell line
HeLa

Genotype
Knockout

Background

This gene encodes a member of the the SLC39A family of divalent metal transporters that mediates the cellular uptake of manganese, zinc, iron, and cadmium. The encoded protein contains eight transmembrane domains, a histidine-rich motif, and a metalloprotease motif, and is expressed on the plasma membrane and the endocytic vesicle membrane. It is an important transporter of nontransferrin-bound iron and a critical regulator of manganese homeostasis. Naturally occurring mutations in this gene are associated with neurodegeneration with brain iron accumulation and early-onset parkinsonism-dystonia with hypermanganesemia.

Gene Information

Gene Symbol
ZIP14

Species
Human

Gene ID
23516

Swiss Prot
Q15043

Synonyms
HCIN; NET34; ZIP14; cig19; HMNDYT2; LZT-Hs4

Product Information

Description
ZIP14 Knockout cell line is engineered from HeLa cell line with Gene-Editing Technology.
Allele-1:19bp deletion in exon1
Allele-2:100bp deletion in exon1
Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging
1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions
Dry ice

Amount
1~5x10^6 cells/vial.

Storage
Store at -20℃

Protocol
Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37℃ with 5% CO_2 condition.
1. Thaw the vial in 37℃ water bath ,and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37℃ incubator with 5% CO_2.
7. A subcultivation ratio of 1:2-1:4 is recommended.


Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Sequencing data



Genome sequence analysis of PCR products from parental (WT) and ZIP14 knockout (KO) HeLa cells, using sanger sequencing.