Leader in Biomolecular Solutions for Life Science

www.abclonal.com

PRKDC Knockdown 293T Cell Line, Heterozygous

Catalog No.: RM50080

Basic Information

Catalog No.
RM50080

Category
Cell Line

Parental Cell line
293T

Genotype
Knockdown

Background

This gene encodes the catalytic subunit of the DNA-dependent protein kinase (DNA-PK). It functions with the Ku70/Ku80 heterodimer protein in DNA double strand break repair and recombination. The protein encoded is a member of the PI3/PI4-kinase family.

Gene Information

Gene Symbol
PRKDC

Species
Human

Gene ID
5591

Swiss Prot
P78527

Synonyms
HYRC; p350; DNAPK; DNPK1; HYRC1; IMD26; XRCC7; DNAPKc; DNA-PKC; DNA-PKcs

Product Information

Description
PRKDC Knockdown cell line is engineered from 293T cell line with Gene-Editing Technology.
Allele-1:18bp deletion in exon1
Allele-2:32bp deletion in exon1
Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging
1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions
Dry ice

Amount
1~5x10^6 cells/vial.


Storage
Store at -20℃

Protocol
Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37℃ with 5% CO_2 condition.
1. Thaw the vial in 37℃ water bath ,and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37℃ incubator with 5% CO_2.
7. A subcultivation ratio of 1:2-1:4 is recommended.

Contact

400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

Sequencing data



Genome sequence analysis of PCR products from parental (WT) and PRKnockdownC knockdown (KD) 293T cells, using sanger sequencing.